

Formation of Higher-dimensional Topological Black Holes

José Natário

(based on arXiv:0906.3216 with Filipe Mena and Paul Tod)

CAMGSD, Department of Mathematics
Instituto Superior Técnico

Talk at IST, 2009



Outline

- 1 Conventions
 - Einstein manifolds
 - Einstein equations
- 2 Collapse to Black Holes without Gravitational Wave Emission
 - Generalized Kottler solutions
 - Generalized Friedman-Lemaître-Robertson-Walker solutions
 - Matching
 - Global properties
- 3 Collapse to Black Holes with Gravitational Wave Emission
 - The Exterior: Bizoń-Chmaj-Schmidt metric
 - The Interiors
 - The Matching

Outline

- 1 Conventions
 - Einstein manifolds
 - Einstein equations
- 2 Collapse to Black Holes without Gravitational Wave Emission
 - Generalized Kottler solutions
 - Generalized Friedman-Lemaître-Robertson-Walker solutions
 - Matching
 - Global properties
- 3 Collapse to Black Holes with Gravitational Wave Emission
 - The Exterior: Bizoń-Chmaj-Schmidt metric
 - The Interiors
 - The Matching

Outline

- 1 Conventions
 - Einstein manifolds
 - Einstein equations
- 2 Collapse to Black Holes without Gravitational Wave Emission
 - Generalized Kottler solutions
 - Generalized Friedman-Lemaître-Robertson-Walker solutions
 - Matching
 - Global properties
- 3 Collapse to Black Holes with Gravitational Wave Emission
 - The Exterior: Bizoń-Chmaj-Schmidt metric
 - The Interiors
 - The Matching

Outline

1 Conventions

- Einstein manifolds
- Einstein equations

2 Collapse to Black Holes without Gravitational Wave Emission

- Generalized Kottler solutions
- Generalized Friedman-Lemaître-Robertson-Walker solutions
- Matching
- Global properties

3 Collapse to Black Holes with Gravitational Wave Emission

- The Exterior: Bizoń-Chmaj-Schmidt metric
- The Interiors
- The Matching

Einstein manifolds

- $(N, d\sigma^2)$ is a n -dimensional **Einstein manifold** with $Ricci = \lambda d\sigma^2$.
- **Examples:** S^n , T^n , H^n , $\mathbb{C}P^{\frac{n}{2}}$, Taub-NUT, Eguchi-Hanson, Calabi-Yau.
- Can be used to construct $(n + 1)$ -dimensional Einstein metrics

$$d\Sigma^2 = d\rho^2 + (f(\rho))^2 d\sigma^2$$

with $Ricci = k d\Sigma^2$.

- If $k > 0$ then $k = \nu^2 n$, $\lambda = \nu^2(n - 1)$, $f = \sin(\nu\rho)$ for some $\nu > 0$.
If $k = 0$ then $\lambda = n - 1$, $f = \rho$ or $\lambda = 0$, $f = 1$.
- If $k < 0$ then $k = -\nu^2 n$, $\lambda = \nu^2(n - 1)$, $f = \sinh(\nu\rho)$ or $k = -\nu^2 n$, $\lambda = 0$, $f = e^{\pm\nu\rho}$ or
 $k = -\nu^2 n$, $\lambda = -\nu^2(n - 1)$, $f = \cosh(\nu\rho)$ for some $\nu > 0$.



Einstein manifolds

- $(N, d\sigma^2)$ is a n -dimensional **Einstein manifold** with $Ricci = \lambda d\sigma^2$.
- **Examples:** S^n , T^n , H^n , $\mathbb{C}P^{\frac{n}{2}}$, Taub-NUT, Eguchi-Hanson, Calabi-Yau.
- Can be used to construct $(n + 1)$ -dimensional Einstein metrics

$$d\Sigma^2 = d\rho^2 + (f(\rho))^2 d\sigma^2$$

with $Ricci = k d\Sigma^2$.

- If $k > 0$ then $k = \nu^2 n$, $\lambda = \nu^2(n - 1)$, $f = \sin(\nu\rho)$ for some $\nu > 0$.
If $k = 0$ then $\lambda = n - 1$, $f = \rho$ or $\lambda = 0$, $f = 1$.
If $k < 0$ then $k = -\nu^2 n$, $\lambda = \nu^2(n - 1)$, $f = \sinh(\nu\rho)$ or $k = -\nu^2 n$, $\lambda = 0$, $f = e^{\pm\nu\rho}$ or
 $k = -\nu^2 n$, $\lambda = -\nu^2(n - 1)$, $f = \cosh(\nu\rho)$ for some $\nu > 0$.



Einstein manifolds

- $(N, d\sigma^2)$ is a n -dimensional **Einstein manifold** with $Ricci = \lambda d\sigma^2$.
- **Examples:** S^n , T^n , H^n , $\mathbb{C}P^{\frac{n}{2}}$, Taub-NUT, Eguchi-Hanson, Calabi-Yau.
- Can be used to construct $(n + 1)$ -dimensional Einstein metrics

$$d\Sigma^2 = d\rho^2 + (f(\rho))^2 d\sigma^2$$

with $Ricci = k d\Sigma^2$.

- If $k > 0$ then $k = \nu^2 n$, $\lambda = \nu^2(n - 1)$, $f = \sin(\nu\rho)$ for some $\nu > 0$.
If $k = 0$ then $\lambda = n - 1$, $f = \rho$ or $\lambda = 0$, $f = 1$.
If $k < 0$ then $k = -\nu^2 n$, $\lambda = \nu^2(n - 1)$, $f = \sinh(\nu\rho)$ or $k = -\nu^2 n$, $\lambda = 0$, $f = e^{\pm\nu\rho}$ or
 $k = -\nu^2 n$, $\lambda = -\nu^2(n - 1)$, $f = \cosh(\nu\rho)$ for some $\nu > 0$.



Einstein manifolds

- $(N, d\sigma^2)$ is a n -dimensional **Einstein manifold** with $Ricci = \lambda d\sigma^2$.
- **Examples:** S^n , T^n , H^n , $\mathbb{C}P^{\frac{n}{2}}$, Taub-NUT, Eguchi-Hanson, Calabi-Yau.
- Can be used to construct $(n + 1)$ -dimensional Einstein metrics

$$d\Sigma^2 = d\rho^2 + (f(\rho))^2 d\sigma^2$$

with $Ricci = k d\Sigma^2$.

- If $k > 0$ then $k = \nu^2 n$, $\lambda = \nu^2(n - 1)$, $f = \sin(\nu\rho)$ for some $\nu > 0$.
If $k = 0$ then $\lambda = n - 1$, $f = \rho$ or $\lambda = 0$, $f = 1$.
If $k < 0$ then $k = -\nu^2 n$, $\lambda = \nu^2(n - 1)$, $f = \sinh(\nu\rho)$ or $k = -\nu^2 n$, $\lambda = 0$, $f = e^{\pm\nu\rho}$ or
 $k = -\nu^2 n$, $\lambda = -\nu^2(n - 1)$, $f = \cosh(\nu\rho)$ for some $\nu > 0$.



Einstein manifolds

- $d\Sigma^2$ typically has a **singularity** at $\rho = 0$: the Kretschmann scalar K is related to the square C^2 of the Weyl tensor of $d\sigma^2$ by

$$K = \frac{C^2}{f^4} + \text{const.}$$

- This can be avoided for $k < 0$ with $f = e^{\pm\nu\rho}$, when the metric has an internal infinity (**cusp**), or $f = \cosh(\nu\rho)$, when the metric has a minimal surface and a second asymptotic region (**wormhole**).



Einstein manifolds

- $d\Sigma^2$ typically has a **singularity** at $\rho = 0$: the Kretschmann scalar K is related to the square C^2 of the Weyl tensor of $d\sigma^2$ by

$$K = \frac{C^2}{f^4} + \text{const.}$$

- This can be avoided for $k < 0$ with $f = e^{\pm\nu\rho}$, when the metric has an internal infinity (**cusp**), or $f = \cosh(\nu\rho)$, when the metric has a minimal surface and a second asymptotic region (**wormhole**).



Einstein equations

- We consider the $(n + 2)$ -dimensional **Einstein equations**

$$R_{ab} = \Lambda g_{ab} + \kappa \left(T_{ab} - \frac{1}{n} T g_{ab} \right),$$

or

$$R_{ab} - \frac{1}{2} R g_{ab} + \frac{n\Lambda}{2} g_{ab} = \kappa T_{ab}.$$



Outline

- 1 Conventions
 - Einstein manifolds
 - Einstein equations
- 2 Collapse to Black Holes without Gravitational Wave Emission
 - Generalized Kottler solutions
 - Generalized Friedman-Lemaître-Robertson-Walker solutions
 - Matching
 - Global properties
- 3 Collapse to Black Holes with Gravitational Wave Emission
 - The Exterior: Bizoń-Chmaj-Schmidt metric
 - The Interiors
 - The Matching

Generalized Kottler solutions

- The **generalized Kottler solutions** are vacuum ($T_{ab} = 0$) solutions given by

$$ds^2 = -V(r)dt^2 + (V(r))^{-1}dr^2 + r^2d\sigma^2,$$

where

$$V(r) = \frac{\lambda}{n-1} - \frac{2m}{r^{n-1}} - \frac{\Lambda r^2}{n+1}.$$



Generalized Friedman-Lemaître-Robertson-Walker solutions

- The Generalized Friedman-Lemaître-Robertson-Walker solutions are dust ($T_{ab} = \mu u_a u_b$) solutions given by

$$ds^2 = -d\tau^2 + R^2(\tau) d\Sigma^2,$$

where $d\Sigma^2$ is any $(n+1)$ -dimensional Riemannian Einstein metric with $Ricci = k d\Sigma^2$ and $R(\tau)$, $\mu(\tau)$ satisfy the conservation equation

$$\mu R^{n+1} = \mu_0$$

and the generalized Friedman equation

$$\frac{\dot{R}^2}{R^2} + \frac{k}{nR^2} = \frac{2\kappa\mu}{n(n+1)} + \frac{\Lambda}{n+1}.$$



Matching

- Can match generalized Kottler to generalized FLRW at $\rho = \rho_0$ provided that $f'(\rho_0) > 0$ and

$$m = \frac{\kappa\mu_0(f(\rho_0))^{n+1}}{n(n+1)}.$$

- Similar results for **generalized Lemaître-Tolman-Bondi**.



Matching

- Can match generalized Kottler to generalized FLRW at $\rho = \rho_0$ provided that $f'(\rho_0) > 0$ and

$$m = \frac{\kappa\mu_0(f(\rho_0))^{n+1}}{n(n+1)}.$$

- Similar results for **generalized Lemaître-Tolman-Bondi**.



Global properties

- If $\Lambda = 0$ (hence $\lambda > 0$) and $(N, d\sigma^2)$ is not an n -sphere then the locally naked singularity is always visible from \mathcal{I}^+ for $k \leq 0$, but can be hidden if $k > 0$ and $n \geq 4$ (cf. [Ghosh and Beesham]).




Figure 1: Penrose diagram for $\Lambda = 0$ and (a) $k \leq 0$; (b) $k > 0$, showing the matching surfaces and the horizons.

Global properties

- If $\Lambda > 0$ (hence $\lambda > 0$) and $(N, d\sigma^2)$ is not an n -sphere then the locally naked singularity can be always be hidden except if the FLRW universe is recollapsing (hence $k > 0$) and $n < 4$.




Figure 2: Penrose diagram for $\Lambda > 0$ with the FLRW universe (a) recollapsing; (b) non-recollapsing, showing the matching surfaces and the horizons.



Global properties

- For $\Lambda < 0$ we have:

If $\lambda > 0$ and $(N, d\sigma^2)$ is not an n -sphere then the locally naked singularity can always be hidden.

If $\lambda = 0$ then the cusp singularity is not locally naked.

If $\lambda < 0$ then no causal curve can cross the wormhole from one \mathcal{I} to the other (cf. [Galloway]).




Figure 3: Penrose diagram for $\Lambda < 0$ and (a) $\lambda > 0$; (b) $\lambda = 0$; (c) $\lambda < 0$, showing the matching surfaces and the horizons.



Outline

- 1 Conventions
 - Einstein manifolds
 - Einstein equations
- 2 Collapse to Black Holes without Gravitational Wave Emission
 - Generalized Kottler solutions
 - Generalized Friedman-Lemaître-Robertson-Walker solutions
 - Matching
 - Global properties
- 3 Collapse to Black Holes with Gravitational Wave Emission
 - The Exterior: Bizoń-Chmaj-Schmidt metric
 - The Interiors
 - The Matching

The Exterior: Bizoń-Chmaj-Schmidt metric

- The **Bizoń-Chmaj-Schmidt metric** is a vacuum ($T_{ab} = 0$) solution of the Einstein equations with $\Lambda = 0$ given by

$$ds^{2+} = -Ae^{-2\delta} dt^2 + A^{-1} dr^2 + \frac{r^2}{4} e^{2B} (\sigma_1^2 + \sigma_2^2) + \frac{r^2}{4} e^{-4B} \sigma_3^2,$$

where σ_i are the standard left-invariant 1-forms on S^3 and A, δ and B are functions of t and r satisfying

$$\partial_r A = -\frac{2A}{r} + \frac{1}{3r} (8e^{-2B} - 2e^{-8B}) - 2r(e^{2\delta} A^{-1} (\partial_t B)^2 + A (\partial_r B)^2);$$

$$\partial_t A = -4rA(\partial_t B)(\partial_r B);$$

$$\partial_r \delta = -2r(e^{2\delta} A^{-2} (\partial_t B)^2 + (\partial_r B)^2);$$

$$\partial_t (e^\delta A^{-1} r^3 (\partial_t B)) - \partial_r (e^{-\delta} A r^3 (\partial_r B)) + \frac{4}{3} e^{-\delta} r (e^{-2B} - e^{-8B}) = 0.$$



The Interiors

- Eguchi-Hanson (S^3/\mathbb{Z}_2):

$$d\Sigma^2 = \left(1 - \frac{a^4}{\rho^4}\right)^{-1} d\rho^2 + \frac{\rho^2}{4}(\sigma_1^2 + \sigma_2^2) + \frac{\rho^2}{4} \left(1 - \frac{a^4}{\rho^4}\right) \sigma_3^2.$$

- k -Eguchi-Hanson (S^3/\mathbb{Z}_p):

$$d\Sigma^2 = \Delta^{-1} d\rho^2 + \frac{\rho^2}{4}(\sigma_1^2 + \sigma_2^2) + \frac{\rho^2}{4} \Delta \sigma_3^2,$$

where $k < 0$, $p \geq 3$, $\Delta = 1 - \frac{a^4}{\rho^4} - \frac{k}{6}\rho^2$ and $a^4 = \frac{4}{3k^2}(p-2)^2(p+1)$.

- k -Taub-NUT:

$$d\Sigma^2 = \frac{1}{4}\Sigma^{-1} d\rho^2 + \frac{1}{4}(\rho^2 - L^2)(\sigma_1^2 + \sigma_2^2) + L^2\Sigma\sigma_3^2,$$

where $\Sigma = \frac{(\rho-L)(1-\frac{k}{12}(\rho-L)(\rho+3L))}{\rho+L}$.



The Interiors

- **Eguchi-Hanson** (S^3/\mathbb{Z}_2):

$$d\Sigma^2 = \left(1 - \frac{a^4}{\rho^4}\right)^{-1} d\rho^2 + \frac{\rho^2}{4}(\sigma_1^2 + \sigma_2^2) + \frac{\rho^2}{4} \left(1 - \frac{a^4}{\rho^4}\right) \sigma_3^2.$$

- **k -Eguchi-Hanson** (S^3/\mathbb{Z}_p):

$$d\Sigma^2 = \Delta^{-1} d\rho^2 + \frac{\rho^2}{4}(\sigma_1^2 + \sigma_2^2) + \frac{\rho^2}{4} \Delta \sigma_3^2,$$

where $k < 0$, $p \geq 3$, $\Delta = 1 - \frac{a^4}{\rho^4} - \frac{k}{6}\rho^2$ and $a^4 = \frac{4}{3k^2}(p-2)^2(p+1)$.

- **k -Taub-NUT**:

$$d\Sigma^2 = \frac{1}{4}\Sigma^{-1} d\rho^2 + \frac{1}{4}(\rho^2 - L^2)(\sigma_1^2 + \sigma_2^2) + L^2\Sigma\sigma_3^2,$$

where $\Sigma = \frac{(\rho-L)(1-\frac{k}{12}(\rho-L)(\rho+3L))}{\rho+L}$.



The Interiors

- **Eguchi-Hanson** (S^3/\mathbb{Z}_2):

$$d\Sigma^2 = \left(1 - \frac{a^4}{\rho^4}\right)^{-1} d\rho^2 + \frac{\rho^2}{4}(\sigma_1^2 + \sigma_2^2) + \frac{\rho^2}{4} \left(1 - \frac{a^4}{\rho^4}\right) \sigma_3^2.$$

- **k -Eguchi-Hanson** (S^3/\mathbb{Z}_p):

$$d\Sigma^2 = \Delta^{-1} d\rho^2 + \frac{\rho^2}{4}(\sigma_1^2 + \sigma_2^2) + \frac{\rho^2}{4} \Delta \sigma_3^2,$$

where $k < 0$, $p \geq 3$, $\Delta = 1 - \frac{a^4}{\rho^4} - \frac{k}{6}\rho^2$ and $a^4 = \frac{4}{3k^2}(\rho - 2)^2(\rho + 1)$.

- **k -Taub-NUT**:

$$d\Sigma^2 = \frac{1}{4}\Sigma^{-1} d\rho^2 + \frac{1}{4}(\rho^2 - L^2)(\sigma_1^2 + \sigma_2^2) + L^2\Sigma\sigma_3^2,$$

where $\Sigma = \frac{(\rho-L)(1-\frac{k}{12}(\rho-L)(\rho+3L))}{\rho+L}$.



The Matching

- In each case, the interior metric gives consistent data for the Bizoń-Chmaj-Schmidt metric at a comoving timelike hypersurface. **Local existence** of the radiating exterior in the neighbourhood of the matching surface is then guaranteed. In the case of **Eguchi-Hanson** and **k -Taub-NUT with $k < 0$** , the data can be chosen to be close to the data for the Schwarzschild solution. Since this solution is known to be stable [**Dafermos and Holzegel**], it is reasonable to expect that the exterior will settle down to the Schwarzschild solution.



Bibliography



Filipe Mena, JN and Paul Tod,
Formation of Higher-dimensional Topological Black Holes,
Annales Henri Poincaré (in press),
arXiv:0906.3216 [gr-qc].



S. Ghosh and A. Beesham,
*Higher-dimensional Inhomogeneous Dust Collapse and Cosmic
Censorship*,
Physical Review D **64** (2001) 124005,
arXiv:gr-qc/0108011.



G. Galloway,
A 'Finite Infinity' Version of Topological Censorship,
Classical and Quantum Gravity **13** (1996) 1471-78.

Bibliography

-  Filipe Mena, JN and Paul Tod,
Formation of Higher-dimensional Topological Black Holes,
Annales Henri Poincaré (in press),
[arXiv:0906.3216](https://arxiv.org/abs/0906.3216) [gr-qc].
-  S. Ghosh and A. Beesham,
Higher-dimensional Inhomogeneous Dust Collapse and Cosmic Censorship,
Physical Review **D 64** (2001) 124005,
[arXiv:gr-qc/0108011](https://arxiv.org/abs/gr-qc/0108011).
-  G. Galloway,
A 'Finite Infinity' Version of Topological Censorship,
Classical and Quantum Gravity **13** (1996) 1471-78.

Bibliography

-  Filipe Mena, JN and Paul Tod,
Formation of Higher-dimensional Topological Black Holes,
Annales Henri Poincaré (in press),
arXiv:0906.3216 [gr-qc].
-  S. Ghosh and A. Beesham,
Higher-dimensional Inhomogeneous Dust Collapse and Cosmic Censorship,
Physical Review **D 64** (2001) 124005,
arXiv:gr-qc/0108011.
-  G. Galloway,
A 'Finite Infinity' Version of Topological Censorship,
Classical and Quantum Gravity **13** (1996) 1471-78.



Bibliography



P. Bizoń, T. Chmaj and B. Schmidt,
Critical Behaviour in Vacuum Gravitational Collapse in $4 + 1$ dimensions,
Physical Review Letters **95** (2005) 071102,
[arXiv:gr-qc/0506074](https://arxiv.org/abs/gr-qc/0506074).



H. Pederson,
Eguchi-Hanson Metrics With a Cosmological Constant,
Classical and Quantum Gravity **2** (1985) 579-587.



M. Dafermos and G. Holzegel,
On the Nonlinear Stability of Higher Dimensional Triaxial Bianchi-IX Black Holes,
Advances in Theoretical and Mathematical Physics **10** (2006) 503-523,
[arXiv:gr-qc/0510051](https://arxiv.org/abs/gr-qc/0510051).



Bibliography



P. Bizoń, T. Chmaj and B. Schmidt,
Critical Behaviour in Vacuum Gravitational Collapse in $4 + 1$ dimensions,
Physical Review Letters **95** (2005) 071102,
[arXiv:gr-qc/0506074](https://arxiv.org/abs/gr-qc/0506074).



H. Pederson,
Eguchi-Hanson Metrics With a Cosmological Constant,
Classical and Quantum Gravity **2** (1985) 579-587.



M. Dafermos and G. Holzegel,
On the Nonlinear Stability of Higher Dimensional Triaxial Bianchi-IX Black Holes,
Advances in Theoretical and Mathematical Physics **10** (2006) 503-523,
[arXiv:gr-qc/0510051](https://arxiv.org/abs/gr-qc/0510051).



Bibliography



P. Bizoń, T. Chmaj and B. Schmidt,
Critical Behaviour in Vacuum Gravitational Collapse in $4 + 1$ dimensions,
Physical Review Letters **95** (2005) 071102,
[arXiv:gr-qc/0506074](https://arxiv.org/abs/gr-qc/0506074).



H. Pederson,
Eguchi-Hanson Metrics With a Cosmological Constant,
Classical and Quantum Gravity **2** (1985) 579-587.



M. Dafermos and G. Holzegel,
On the Nonlinear Stability of Higher Dimensional Triaxial Bianchi-IX Black Holes,
Advances in Theoretical and Mathematical Physics **10** (2006) 503-523,
[arXiv:gr-qc/0510051](https://arxiv.org/abs/gr-qc/0510051).

